

Blue Variable Stars from the MACHO database I: Photometry and Spectroscopy of the LMC sample

S. C. Keller

Institute of Geophysics and Planetary Physics, LLNL,
Livermore, CA 94550

`skeller@igpp.ucllnl.org`

M. S. Bessell

Research School of Astronomy and Astrophysics, ANU,
Canberra, Australia

K. H. Cook

Institute of Geophysics and Planetary Physics, LLNL,
Livermore, CA 94550

M. Geha

Dept. of Astronomy and Astrophysics,
UCO/Lick Observatory, Univ. of California, Santa Cruz, CA 95064

and

D. Syphers

Center for Cosmological Physics, University of Chiago,
5640 S. Ellis Ave., Chicago, IL 60637

Received _____; accepted _____

Submitted to Astronomical Journal

ABSTRACT

We present the photometric properties of 1279 blue variable stars within the LMC. Photometry is derived from the MACHO database. The lightcurves of the sample exhibit a variety of quasi-periodic and aperiodic outburst behavior. A characteristic feature of the photometric variation is that the objects are reddest when at maximum outburst. A subset of 102 objects were examined spectroscopically. Within this subset, 91% exhibited Balmer emission in at least one epoch, in some cases with spectacular spectral variability. The variability observed in the sample is consistent with the establishment and maintenance of the Be phenomenon.

Subject headings: stars: emission-line, Be — stars: variables: other — galaxies: individual (Large Magellanic Cloud)

1. Introduction

The MACHO photometric database constitutes an unprecedented resource for the study of stellar variability. With coverage of over 20 million stars within the LMC alone, the MACHO project is superior to competitors in duration (seven years) and spatial coverage. A primary goal of the MACHO project was to detect microlensing events in/towards the Magellanic Clouds and the bulge of the Galaxy. One of the initial unexpected contaminants in the detection of microlensing events were blue variables provisionally termed “bumpers” (Cook et al. 1995). These objects were found to brighten $\Delta V \sim 0.^m2\text{--}0.4$ over the course of fifty to several hundred days in a manner much like that expected from microlensing events. However these objects did not exhibit the achromaticity expected of microlensing.

In the present work we extract from the catalogue of MACHO variables (over

210000 objects) those stars which are in the vicinity of the main-sequence and exhibit large amplitude variability (see Section 2). Section 3 presents an overview of lightcurve morphology and colour variations. Section 4 presents spectroscopic observations of a subset of 102 objects from our catalogue. In Section 5 we discuss the physical nature of the blue variables. The goal of the current paper is to present preliminary data on these objects for future studies to expand upon.

2. Selection of the Sample

Instrumental MACHO B and R magnitudes have been transformed to standard Johnson V and Cousins R (for details see Alcock et al. 1999). From this database we have selected blue (i.e. $-0.5 < V - R < 0.4$) objects with $V < 18.0$. We then select those stars with significant variability. Our variability index borrows from that used to detect MACHO microlensing events (Alcock et al. 1995). Under this scheme, a star is flagged as variable if the lightcurve contains at least 10 pairs of V and R of large χ^2 compared to a constant V and R brightness. From the 30 LMC MACHO fields with calibrated photometry this results in a sample of 2841 candidate variables. A major contaminant are eclipsing binaries which were excluded by visual inspection. This results in a sample of 1279 objects. The positions, magnitudes and colours of these objects are presented in Table 2 (full table is available from ADS).

3. Photometry

Our selected blue variables show a remarkable variety of lightcurve morphology. To attempt to classify each star into a series of arbitrary categories is perhaps misleading as stars may show combinations of what might be termed “modes” of variability. These

“modes” can be described as follows:

1. Bumper events - these events are typically $\Delta V = 0.^m2-0.^m4$ in amplitude and duration of 100-800 days.
2. Flicker events - rapid, generally low amplitude ($\Delta V = 0.^m05-0.^m15$) variation. Duration of outburst ranges from 10-50 days. Rise time is very short (several days) followed by “exponential” decay. Larger outbursts result in longer durations. Flickering behavior is often associated with maximum light of bumper modes (Fig. 2).
3. Step events - within the course of 10-50 days the brightness increases $\Delta V = 0.^m2-0.^m3$.
4. Baseline variation - long-term trend of 1000s of days with amplitudes up to $\Delta V \sim 0.^m4$.
5. Fading events - drop in V luminosity of up to $\Delta V \sim 0.^m4$ with time scales 200-600 days.

Figure 1 shows examples of these modes. As can be seen from Figure 2 many stars exhibit modes 1-5 in combination. The $V-R$ colours show that outburst events result in redder colours. Whilst modes 1,2 and 5 bring about a simultaneous change in colour, modes 3 and 4 often exhibit a long delay ($\sim 1000d$) between attaining maximum light and attaining reddest colour.

The seven year baseline of the MACHO photometry shows the ephemeral nature of each mode. A particular mode of behavior may switch on or off without precursor. For this reason we propose that the five modes of variability are expressions of one underlying mechanism which we discuss below. In a small number of our sample (324 from 1279 stars) the variability appears quasi-periodic albeit with variable amplitude. Amongst quasi-periodic stars, periods range from 150-1400d for mode 1 with a mean period of 560d to 20-100d for objects of mode 2 (mean $P=65d$).

Figure 4 shows the V , $V-R$ colour-magnitude diagram (CMD) for the sample of blue variables we have isolated. These stars form a broad band parallel to the non-variable stars on the main-sequence. This red offset is of similar magnitude to that observed amongst the Magellanic Cloud Be star population (Keller, Wood & Bessell 1999). Shown in large symbols are those objects which exhibit only one mode in the available data. The five modes inhabit a similar range in colour. Figure 5 presents the luminosity functions for the subset of single mode stars. At high luminosity, large outburst behaviour (mode 1) dominates the blue variable population. The smaller mode 2 events are significantly skewed to lower luminosities.

4. Spectroscopy

Spectroscopy was performed on a subset of 102 blue variables. These observations were conducted on eight epochs however only 39 stars have two or more observations. Table 1 describes our spectroscopic observations. The sample of objects observed spectroscopically contains stars from each of the five modes discussed above.

Amongst the stars in our sample, 91% show Balmer line emission in at least one epoch. That is, the majority of objects are Be stars. Given the fact that such a high proportion of the sample show emission in our limited number of epochs and the variable nature of the emission-line phenomenon it is highly likely that all stars in our sample are Be stars. Amongst the 39 stars with multiple spectroscopic observations $\sim 50\%$ show dramatic variation in emission equivalent width of H Balmer lines. The most variable are those which undergo outburst from a quiescent constant baseline. An example of this extreme variability is shown for three stars in Fig. 3. The $H\alpha$ line profile is seen to vary from absorption to equivalent widths of over $-60\text{\AA}$. Time of maximum light is associated with maximum Balmer line emission in each instance. Those which do not exhibit a constant

baseline are characterised by strong emission (typically -20 - $-40\text{\AA}$).

5. The Nature of Blue Variables

The blue variables of the LMC exhibit a variety of lightcurve morphology yet present similar colour, luminosity distribution and spectral type. We believe that the lack of differences between objects exhibiting different modes is due to the fact that the variability has a common underlying mechanism. The Be phenomenon, and the process underlying it, is the prime suspect for the production of the variability. A Be star is defined simply as a B star that has at one time exhibited Balmer emission. We find that 91% of the blue variable population examined spectroscopically are Be stars.

An early reference to the unusual population of blue variables in the LMC was made by Hodge (1961) who discussed a class of variable star that matches that outlined here, namely, stars to the red of the main-sequence presenting eruptive outbursts which are redder when brighter. Subsequently, the EROS microlensing survey have also discussed the blue variable population in the LMC (Beaulieu et al. 1996; Lamers, Beaulieu & de Wit 1999). These studies highlighted a small sample of seven objects which satisfy selection criteria similar to those applied here. These studies claim however, that the blue variables belong to a pre-main-sequence (PMS) population analogous to Galactic Herbig Ae/Be stars. Study of the EROS SMC data by Beaulieu et al. (2001) and de Wit, Beaulieu & Lamers (2002) reveals a further seven similar objects.

The five of the seven EROS LMC objects are present within our catalogue of blue variables. The counterparts of two objects are not present because they do not exhibit significant variability in our dataset. Seen in the context of our eight year baseline rather than the limited 117 days of EROS, these objects do not show variability which sets them

apart from the remainder of our sample.

Figure 6 presents the lightcurves for ELHC1-3 & 6 (ELHC5 was observed in MACHO R band only but is clearly of mode 1). The variability exhibited is distinct from the frequent, large amplitude (up to 3 mag.) fading events of short duration (2-50 days) seen in Galactic Herbig Ae/Be stars (Rostopchina & Grinin 2001; Bibo & The 1991; The 1994). To definitively distinguish between the possible Be and Herbig Ae/Be nature of these objects will require further study particularly infra-red spectroscopy.

We propose that the variability of these stars is the result of processes related to the establishment, maintenance and dissipation of the Be disk. The emission that characterises Be stars originates in a gaseous circumstellar quasi-Keplerian disk. No definitive model has been developed which can sufficiently describe the formation and maintenance of this disk material. The wind compressed disk model (Bjorkman & Cassinelli 1993) utilises the combination of fast photospheric rotation with large mass loss rates to generate a confined disk of material but falls short of generating the densities observed within Be star disks. Other models such as the “one-armed” density wave model (Okazaki 1997) seek to explain spectral line variations observed in Be stars.

An insight into the establishment of Be disks is provided by the works of Rivinius et al. (1998; 2001) in their detailed study of the Be star μ Cen. These authors find evidence for non-radial pulsation modes within the star which when in constructive interference provide sufficient mechanical work to expel material from the stellar surface. Such material gives rise to short lived spectral variations which damp as the material melds into the bulk of the Keplerian disk. The outbursts on μ Cen have remained of low intensity and result in only minor modulation of the V luminosity.

A large proportion of galactic Be stars are known to exhibit photometric and spectroscopic variability. Many studies exist in the literature which have sought to

characterise the nature of these variations (Feinstein & Marraco 1979; Dachs 1982; Hubert & Floquet 1998). The study by Hubert & Floquet of Hipparcos photometry of 289 Be stars shows that amongst early type Be stars variability greater than $0.^m02$ (in the broadband Hipparcos photometric system) is commonplace (98% amongst B0-B3 stars). Hubert & Floquet describe five types of variability which encompass those defined for the present sample. Short-term variations (hours to days) are low amplitude (a few hundredth of a mag.) and are frequently periodic in nature. Mid-term variations (days to weeks) are seen to occur as outbursts of several tenths of a magnitude which are both short-lived (~ 50 days: e.g. ω CMa; like mode 2) and long-lived (~ 500 days: e.g. ν Cyg; like mode 1) and in the form of fading events. Long-term variations (years) of upto $0.^m12$ are seen over the 1200d Hipparcos timespan.

The photometric behaviour of Galactic Be stars as described by Hubert & Floquet mimics that seen amongst the present sample. In particular the bumper and flicker outburst modes are of similar duration and amplitude. Whilst the present sample of blue variables is undoubtedly heterogeneous we believe that the variability of the majority of objects can be best understood in connection with the Be phenomenon.

Let us now interpret our observations in terms of the Be phenomenon. Our spectroscopy indicates that objects exhibiting mode 1 or 2 outbursts from a flat or quiescent baseline exhibit maximum line emission when in outburst and an absence of emission when in quiescence. Each outburst event can be clearly interpreted as a mass-ejection event from the stellar surface. The ejection of material into the circumstellar environment creates a rise in luminosity and redder colour due to the addition of Balmer continuum emission from the cooler disk component (Zorec & Briot 1991). A slow decline follows the initial outburst as material is removed from the circumstellar environment due to outflow (see Rivinius et al. 2001). In this proposed scenario, mode 1 and 2 outbursts arise from the same mechanism,

however, in mode 1 more substantial mass loss occurs. Outbursts may also occur in systems with established circumstellar material leading to the variety of lightcurve morphology seen in Fig. 2. Similarly, fading events (mode 5) reflect the reduction of circumstellar material. Unfortunately we do not have multi-epoch spectroscopic observations over a fading event to confirm this.

The blue variables we have presented in this work afford an excellent probe into the mechanisms of establishment and maintenance of the Be phenomenon. Further study of the combined photometric and spectroscopic variations may infer the mechanisms responsible for mass ejection and establishment of the circumstellar disk.

6. Summary

We have presented a summary of the properties of 1279 blue variables within the Large Magellanic Cloud. Variations of up to $\Delta V \sim 0.^m4$ are observed in the form of rapid (~ 50 d) or prolonged (100-500d) eruptive events as well as fading events and long term (years) trends. The $V-R$ colours show a characteristic “redder when brighter” behaviour.

A subset of 102 stars were examined spectroscopically, in a limited number of instances over multiple epochs. Many objects show spectacular photometric and spectroscopic variability. The majority (91%) of the spectroscopic sample were found to exhibit Balmer emission lines in at least one epoch (i.e. Be stars).

This strongly suggests that the mechanism giving rise to the variability seen with in the sample of LMC stars is related to the Be phenomenon. Firstly, the photometric variability is similar to that observed in Galactic Be stars. Secondly, emission strength is seen to coincide with redder $V-R$ colour and maximum V luminosity as expected from known properties of Galactic Be stars.

The outbursts observed in the blue variable population can therefore be understood as mass eject events from the central star leading to the establishment and maintenance of the Be disk. Through further study of this population could yield important insight into the physical process of mass loss and the confinement of material in the vicinity of star which underlies the Be star phenomenon. The role of metallicity in the Be phenomenon will be examined in a future paper where we plan to investigate properties of the blue variable population of the Small Magellanic Cloud.

This work was performed under the auspices of the U. S. Department of Energy, National Nuclear Security Administration by the University of California, Lawrence Livermore National Laboratory under contract W7405-Emg-48.

REFERENCES

- Alcock, C. et al. 1995, AJ, 109, 1653
- Alcock, C. et al. 1999, PASP, 111, 1539
- Balona, L.A., & Shobbrook, R.R. 1984, MNRAS, 211, 375
- Beaulieu, J. P. et al. 1996, Science, 272, 995
- Beaulieu, J.-P. et al. 2001, A&A, 380, 168
- Bibo, E.A., & The, P.S. 1991, A&AS, 89, 319
- Bjorkman, J. E. & Cassinelli, J. P. 1993, ApJ, 409, 429
- Cook, K. H. et al. 1995, in Astrophysical Applications of Stellar Pulsation, ASP Conf. Ser. 83, ed. R. S. Stobie & P. A. Whitelock (San Francisco:ASP), 221
- de Wit, W.J.M., Beaulieu, J-P. & Lamers, H. A&A, submitted
- Dachs, J. 1982, in IAU Symp. 98, Be stars, ed. M. Jaschek & H.-G. Groth (Dordrecht: Reidel), 19
- Feinstein, A., & Marraco, H.G. 1979, AJ, 84, 1713
- Hodge, P.W. 1961, Obs 81, 31
- Hubert, A.M., & Floquet, M. A&A, 335, 565
- Keller, S. C., Wood, P. R., & Bessell, M. S. 1999, A&AS, 134, 489
- Lamers, H. J. G. L. M., Beaulieu, J. P., & de Wit, W. J. 1999, A&A, 341, 827
- Okazaki, A. T. 1997, A&A, 318, 548

Rivinius, T., Baade, D., Štefl, S., Stahl, O., Wolf, B., & Kaufer, A. 1998, *A&A*, 333, 125

Rivinius, T., Baade, D., Štefl, S., & Maintz, M. 2001, *A&A*, 379, 257

Rostopchina, A. N. & Grinin, V. P. 2001, *Astronomy Reports*, 45, 51

The, P. S. 1994, in *ASP Conf. Ser. 62: The Nature and Evolutionary Status of Herbig Ae/Be Stars*, ed. P. S. The, M. R. Perez, and E. P. J. Van den Heuvel (San Francisco: ASP), 23

Zorec, J. & Briot, D. 1991, *A&A*, 245, 150




Fig. 1.— V lightcurves and $V-R$ colourcurves for examples of the five modes of variability exhibited by the sample of blue variables. The MACHO identifications from top to bottom are 17.2109.68, 77.7672.34, 14.8500.382, 81.9490.41, and 1.4050.1666.




Fig. 2.— Example V lightcurves and $V-R$ colourcurves for objects showing the variety of behaviour obtained from juxtaposition of the five modes of variability defined in Fig. 1. The MACHO identifications from top to bottom are 10.3195.13, 1.3813.48, 10.4036.11, 17.2229.18, and 1.3931.95




Fig. 3.— Three objects with multiple spectroscopic observations (Top:79.4780.145, Middle:6.6571.71, Bottom:78.6223.41). Note the extreme spectra variability between the two epochs indicated by the dotted vertical lines on the lightcurve. In each case the epoch of maximum emission coincides with that of maximum light.




Fig. 4.— The $V, V-R$ colour-magnitude diagram for the blue variables described in the present work. Symbols are as follows: mode 1 - solid squares, mode 2 - stars, mode 3 - open squares, mode 4 - asterisk and mode 5 - solid circles. The zero-age main-sequence is shown for comparison (Balona & Shobbrook 1984)




Fig. 5.— Luminosity functions of those stars exhibiting only one mode of variability over the duration of our monitoring. Mode 1 is shown dashed, mode 2 solid, mode 4 dotted and




Fig. 6.— V lightcurves and $V-R$ colourcurves for four pre-main sequence candidates identified by Lamers, Beaulieu & de Wit (1999) (MACHO identifications are 78.6101.55(top), 78.6220.76, 79.5863.73, 78.6100.101). Our eight year baseline shows that these objects do not presently variations of significantly different nature to those presented in Figs. 1 & 2.

Table 1: Details of our spectroscopic observations

date	MJD	Instrument	$\lambda(\text{\AA})$	Resolution $\Delta\lambda(\text{\AA})$
30 Dec 1994	49717	AAT/RGO	6290-6840	1.0
2 Jan 1997	50451	CTIO 60"	3640-7330	4.4
26 Nov 1998	51144	SSO 2.3m/DBS	3600-4580; 6095-7050	1.1; 1.1
28 Mar 1999	51266	SSO 2.3m/DBS	3315-5500; 5500-10200	4.4; 8.0
22 Sep 1999	51444	SSO 2.3m/DBS	3250-6410; 6400-10400	4.4; 8.0
12 Dec 1999	51525	SSO 2.3m/DBS	3600-4600; 6100-7050	1.1; 1.1
2 Dec 2001	52246	SSO 2.3m/DBS	3610-4590; 6090-7045	1.1; 1.1
4 Mar 2002	52310	SSO 2.3m/DBS	3610-4590; 6090-7045	1.1; 1.1

Table 2: Positions and mean photometry for the sample of 1279 blue variables. The full table is available in electronic form from ADS. The column “mode” describes the most prominent of the five modes outlined in section 3. In some cases where a number of modes are apparent these are also given. The full table is available from ADS.

MACHO id	$\alpha(\text{J2000})$	$\delta(\text{J2000})$	V	$V - R$	mode
1.3321.7	05:01:13.700	-69:20:04.24	14.756	14.671	1
1.3442.19	05:01:27.791	-69:19:51.16	14.747	14.729	5
1.3446.1061	05:01:35.476	-69:04:28.83	16.564	16.494	4
1.3563.47	05:02:32.764	-69:21:18.40	15.756	15.676	24
1.3564.154	05:02:31.128	-69:14:36.92	17.182	17.048	1
...	...	...	...	...	...